




SOCIEDADE BRASILEIRA
DE ENTOMOLOGIA
FUNDADA EM 1937

REVISTA BRASILEIRA DE
Entomologia
A Journal on Insect Diversity and Evolution

www.rbentomologia.com



Short Communication

First host record of *Epipompilus* (Hymenoptera: Pompilidae) from Brazil and discussion of prey carriage mechanism



Bhrenno M. Trad^{a,b}, Rogerio Silvestre^{a,b,*}, Tiago H. Auko^a, Vinicius M. Lopez^a, Eduardo F. dos Santos^c

^a Universidade Federal da Grande Dourados, Faculdade de Ciências Biológicas e Ambientais, Laboratório de Ecologia de Hymenoptera, Dourados, MS, Brazil

^b Universidade Federal da Grande Dourados, Faculdade de Ciências Biológicas e Ambientais, Programa de Pós-Graduação em Entomologia e Conservação da Biodiversidade, Dourados, MS, Brazil

^c Universidade Estadual Paulista “Júlio de Mesquita Filho”, Instituto de Biociências, Letras e Ciências Exatas, Departamento de Zoologia e Botânica, São José do Rio Preto, SP, Brazil

ARTICLE INFO

Article history:

Received 15 June 2018

Accepted 20 August 2018

Available online 3 September 2018

Associate Editor: James Carpenter

Keywords:

Behavior

Brazilian Cerrado

Host association

Segestriidae

Spider wasp

ABSTRACT

We register for first time the occurrence of *Epipompilus tucumanus* Evans, 1967 in Brazil, and record the spider *Ariadna boliviana* Simon, 1907 as its host. The observations were made in the National Park of Chapada dos Guimarães, Mato Grosso, Brazil. The prey carriage mechanism is described for first time for this genus, and we provide a video showing this behavior.

© 2018 Sociedade Brasileira de Entomologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

The spider wasp genus *Epipompilus* Kohl, 1884 comprises 51 species, 17 of which occur in the Neotropical Region and 34 in the Australian Region (Evans, 1953; Elliot, 2007). The biology of this genus is poorly known, and prey has been reported for only two species: *E. insularis* Kohl, 1884 (Pollard, 1982; Harris, 1987) and *E. platensis* (Roig-Alsina and Barneche, 2017). Six spider families have been reported up to now as *Epipompilus* host: Clubionidae, Desidae, Heteropodidae, Salticidae, Sparassidae and Segestriidae (Evans, 1972; Pollard, 1982; Harris, 1987; Shimizu, 1994; Roig-Alsina and Barneche, 2017).

Evans (1967) describes *Epipompilus tucumanus* from Tucuman and Jujuy (Argentina) in Chacoan biogeographical province, and also from San Esteban, Venezuela, at the Venezuelan biogeographical province. Evans (1976) added another locality from Buena Vista in Santa Cruz, Bolivia, at Rondonian biogeographical province.

On March 28, 2016, around 11 AM, at the National Park of Chapada dos Guimarães, MT, Brazil (−15.4069° −55.8239°, 610m asl.) (Fig. 1), two of us (B.M.T. and V.M.L.) observed a female of *E. tucumanus* (Fig. 2) carrying a cursorial spider on a trail inside the Cerrado *sensu stricto*. According to Morrone (2014) this site is located in the Cerrado biogeographical province, Chacoan sub-region, and Chacoan dominion. The climate of this

region is classified by Köppen into Tropical with dry winter (Aw) (Alvares et al., 2013). The wasp was collected and deposited in the Hymenoptera collection of the Museu de Biodiversidade (MuBio), Universidade Federal da Grande Dourados (UFGD), Mato Grosso do Sul State, Brazil (voucher number Hym-00191-P). The spider was determined by Dr. Antônio D. Brescovit, and deposited in the arachnid's collection of the Instituto Butantan, São Paulo State, Brazil (voucher number IBSP-211845).

We sent pictures of the specimen collected in Chapada dos Guimarães to Dr. Arturo Roig-Alsina for identification, and we compared it with the images of the holotype available at <https://mczbase.mcz.harvard.edu/guid/MCZ:Ent:31320>. We agree that the specimen is a variation of *E. tucumanus* Evans (1967). It differs from Evans's description by the followings characters: 1- the ivory-white spot is present only on the scape (Fig. 3), Evans (1967) comments that such spots are present on the first three antennal segments; 2- the mesoscutum is wholly black, without ferruginous midline posteriorly (Fig. 4); 3- the whitish band on the posterior margin of the pronotum has no rufous border (Figs. 3 and 4); 4- the first gastral segment and the basal 2/3 of tergum II are ferruginous, with the remainder black (Fig. 5); 5- the whitish markings on meso- and hindtibiae are reduced on the outer and the inner surface (Fig. 6); 6- the clypeus is 2.0× as wide as its median length (Fig. 7); 7- the middle interocular distance equals 0.51× width of the head (Fig. 7); 8- the lower interocular distance is equal to 0.9× of the upper one (Fig. 7); 9- unlike the holotype, our specimen has oval

* Corresponding author.

E-mail: rogeriosilvestre@ufgd.edu.br (R. Silvestre).




Fig. 1. Collecting localities of *Epipompilus tucumanus* Evans, 1967 in South America, with biogeographic provinces (Morrone, 2014), including the type locality and Evans (1976) record. Highlighting the Chapada dos Guimarães National Park limits, with our collecting point (red star).



Fig. 2. *E. tucumanus* ♀ lateral habitus.



Fig. 3. *E. tucumanus* ♀ dorsal view of head and thorax.



Fig. 4. *E. tucumanus* ♀ thorax in lateral view.



Fig. 5. *E. tucumanus* ♀ metasoma in dorsolateral view.



Fig. 8. *E. tucumanus* ♀ front ventral view of head.



Fig. 6. *E. tucumanus* ♀ hind tibia.



Fig. 9. *Ariadna boliviana* dorsal view.



Fig. 7. *E. tucumanus* ♀ frontal view of head.



Fig. 10. *Ariadna boliviana* ventral view.

white spots between the antennal lobes and eyes (Fig. 8). In view of these differences possibly the specimen represents an undescribed species.

The spider captured by *E. tucumanus* is an adult female of *Ariadna boliviana* Simon, 1907 (Araneae: Segestriidae) (Figs. 9 and 10), and this is the first prey record for *E. tucumanus*. The genus *Ariadna* (Segestriidae) was recently registered as host of *Epipompilus platenis* from Argentina (Roig-Alsina and Barneche, 2017). This spider is nocturnal, sedentary and usually lives in silk tubes constructed in trunk or rock holes (Capocasale, 1998; Grismado, 2008). *Ariadna* Audouin, 1826 occurs on all continents except Antarctica, and *A. boliviana* is known from Brazil, Bolivia, Suriname and Paraguay (Giroti, 2013).

Epipompilus tucumanus, as other spider wasps, uses the mandibles for prey carrying. The wasp drags the spider walking sideways and sometimes backwards, mostly keeping it by the spinnerets, but also can hold the prey by the legs or chelicera to overcome obstacles in the leaf litter like twigs and stones (Videos 1 and 2).

The prey carriage mechanism is similar in *Dipogon* Fox, 1897 (Pompilidae), in which prey carriage occurs backwards and sideways, also the way these wasps hold their prey (Evans and Yoshimoto, 1962; Kurczewski et al., 2017). Evans (1962) classifies this behavior as a variation of Mandibular Mechanism type One (M1): simply grasp the host with their mandibles on any convenient part of the body and drag it backwards into a hole. Genise (1980) comments that this variant hardly is a transition between different

kinds of prey carriage mechanisms, and that carriage behavior is related to the prey size and weight, proposing an exclusive variant for mandibular carriage mechanism type (Variant 5).

The maternal behavior varies significantly in *Epipompilus*. Pollard (1982) and Harris (1987) observed in New Zealand that *E. insularis* oviposits on the spider's body immediately after having paralyzed it (the paralysis is very light and short); no transportation to a nest was observed. After laying the egg the wasp moves away and the spider retakes its normal activities until the wasp larva kills it. Evans (1953) expressed the sequence of behavior stated for *E. insularis* by the formula VPO (*Venari* = to hunt; *Pungere* = to sting the prey; *Ovum parere* = to lay the egg) or VO, and expressed the opinion that such ethological sequences represents primitive conditions within Aculeata. In contrast, *E. tucumanus* showed a more complex ethological sequence that should be described as VPOTC (*Venari*; *Pungere*; *Transferre* = to carry the prey; *Ovum parere*; *Claudere* = to close the cell); this last formula applies to every species of Pompilidae that carries its host to a nest and encloses it in a cell. Evans (op. cit.) thought that VPOTC is the most primitive behavioral sequence within Pompilidae, and that VPO could be derived from VPOTC.

Conflict of interest

The authors declare no conflicts of interest.

Acknowledgments

We sincerely thanks Dr. Arturo Roig-Alsina for species identification; Dr. Antônio D. Brescovit for spider identification, and Dr. Wojciech Pulawski for review the manuscript; the researchers Isabelli L. T. de Jesus, Larissa T. da Silva and Vander Carbonari for their field assistance; ICMBio's staff from Parque Nacional da Chapada dos Guimarães, for the holding in the field. Collect license number: ICMBio 51296-1. B.M.T. is supported by CAPES, process number: PDSE 88881.131920/2016-01.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.rbe.2018.08.003](https://doi.org/10.1016/j.rbe.2018.08.003).

References

- Alvares, C.A., Stape, J.L., Sentelhas, P.C., Gonçalves, J.L.M., Sparovek, G., 2013. Köppen's climate classification map for Brazil. *Meteorol Z* 22 (6), 711–728.
- Capocasale, R.M., 1998. Arañas del Uruguay, VIII. Contribución al conocimiento de *Segestria ruficeps* Guérin (Araneae, Segestriidae). *Comunicaciones Zoológicas del Museo de Historia Natural de Montevideo* 190, 1–12.
- Elliot, M.G., 2007. Annotated catalogue of the Pompilidae (Hymenoptera) of Australia. *Zootaxa* 1428 (1), 1–83.
- Evans, H.E., 1953. Comparative ethology and the systematics of spider wasps. *Syst Zool* 2, 155–172.
- Evans, H.E., 1962. The evolution of prey-carrying mechanism in wasps. *Evolution* 16, 468–483.
- Evans, H.E., 1967. Studies on neotropical Pompilidae (Hymenoptera) III. Additional notes on *Epipompilus* Kohl. *Breviora* 273, 1–15, Available from: <http://biostor.org/reference/4267>.
- Evans, H.E., 1972. Revision of the Australian and New Guinean species of *Epipompilus* (Hymenoptera: Pompilidae). *Pac Insects* 14, 101–131.
- Evans, H.E., 1976. Studies on Neotropical Pompilidae (Hymenoptera). X. Supplementary notes. *Psyche* 83, 263–270.
- Evans, H.E., Yoshimoto, C.M., 1962. The ecology and nesting behaviour of the Pompilidae of the Northeastern United States. *Misc Publ Ent Soc Amer* 3, 67–119.
- Genise, J.F., 1980. Evolución de mecanismos de transporte de presas en avispa. Reclasificación de los tipos mandibulares y esquema evolutivo general. *Rev Soc Entomol Arg* 39, 19–36.
- Giroti, A.M., (Ph.D. thesis) 2013. Revisão taxonômica das aranhas neotropicais do gênero *Ariadna* Audouin, 1826 (Araneae Segestriidae), Universidade de São Paulo, São Paulo, Brasil, 146pp.
- Grismado, C.J., 2008. A taxonomic revision of the spider genus *Ariadna* Audouin, 1826 in Argentina and Chile, with the description of five new species (Arachnida, Araneae, Segestriidae). *Zoosystema* 30 (2), 333–360.
- Harris, A.C., 1987. Pompilidae (Insecta: Hymenoptera). *Fauna of New Zealand* 12, 1–154.
- Kurczewski, F.E., Edwards, G.B., Pitts, J.P., 2017. Hosts, nesting behavior, and ecology of some North American spider wasps (Hymenoptera: Pompilidae), II. *Southeast Nat* 16 (m9), 1–82.
- Morrone, J.J., 2014. Biogeographical regionalisation of the Neotropical region. *Zootaxa* 3782 (1), 001–110.
- Pollard, S.D., 1982. *Epipompilus insularis* (Hymenoptera: Pompilidae), a parasitoid of hunting spiders. *New Zeal J Zool* 9, 37–39.
- Roig-Alsina, A., Barneche, J., 2017. The genus *Epipompilus* in Argentina (Hymenoptera: Pompilidae). *Rev Soc Entomol* 76 (3–4), 33–38.
- Shimizu, A., 1994. Phylogeny and classification of the family Pompilidae. *Tokyo Metropolitan University Bulletin of Natural History* 2, 1–142.